

BRAIDS, TRANSVERSAL LINKS AND THE KHOVANOV-ROZANSKY THEORY

HAO WU

ABSTRACT. We establish some inequalities about the Khovanov-Rozansky cohomologies of braids. These give new upper bounds of the self-linking numbers of transversal links in standard contact S^3 which is sharper than the well known bound given by the HOMFLY polynomial. We also introduce a sequence of transversal link invariants, and discuss some of their properties.

1. INTRODUCTION

In [7, 8], M. Khovanov and L. Rozansky introduced two versions of Khovanov-Rozansky link cohomologies. Their construction is based on matrix factorizations associated to certain planar diagrams. The version in [7] uses the potential x^{n+1} , which lead to a $\mathbb{Z} \oplus \mathbb{Z}$ -graded cohomology theory H_n for each $n \geq 2$. The version in [8] uses the potential ax , which gives a $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ -graded cohomology theory H .

M. Khovanov and L. Rozansky showed that H_n is invariant under Reidemeister moves, and the isomorphism type of H is invariant under Markov moves of closed braids up to overall shifts of gradings. The graded Euler characteristics of H_n and H are variants of the HOMFLY polynomial.

In this paper, we discuss some applications of these theories in contact topology, specifically the transversal knot theory. The sign convention of this paper is given in figure 1, which is opposite to that used in [7, 8].



FIGURE 1. Sign of a crossing

First, we introduce a variant of H whose graded isomorphism type is invariant under Reidemeister moves. Let B be a closed braid with b strands and writhe w . Define $\hat{H}(B)$ to be $H(B)$ with each of the two quantum gradings shifted by $\frac{w+b}{2}$ and the homological grading shifted by $-\frac{w+b}{2}$. Then $\hat{H}(B)$ is $\frac{1}{2}\mathbb{Z} \oplus \frac{1}{2}\mathbb{Z} \oplus \frac{1}{2}\mathbb{Z}$ -graded. From M. Khovanov and L. Rozansky's construction, it's clear that the gradings of $\hat{H}(B)$ are now invariant under all Markov moves.

2000 *Mathematics Subject Classification.* Primary 57M25, 57R17.

Key words and phrases. Braid, transversal knot, knot homology, matrix factorization.

Denote by $H_n^{j,k}$ the space of homogeneous elements with grading (j, k) of H_n , where j is the homological grading, and k is the quantum grading which comes from the exponents of the markings x_i . Denote by $\hat{H}^{j,k,l}(B)$ the space of homogeneous elements with grading (j, k, l) of $\hat{H}(B)$, where j is the cohomological grading, k is the quantum a -grading which comes from the exponents of the formal variable a , and l is the quantum x -grading which comes from the exponents of markings x_i .

For a link L , define

$$\begin{aligned} g_{\max}^{(n)}(L) &= \max\{k \mid H_n^{j,k}(L) \neq 0 \text{ for some } j\}, \\ g_{\min}^{(n)}(L) &= \min\{k \mid H_n^{j,k}(L) \neq 0 \text{ for some } j\}, \\ \hat{g}_{\max}(L) &= \max\{k \mid \hat{H}^{j,k,l}(L) \neq 0 \text{ for some } j, l\}, \\ \hat{g}_{\min}(L) &= \min\{k \mid \hat{H}^{j,k,l}(L) \neq 0 \text{ for some } j, l\}. \end{aligned}$$

These are numerical link invariants. We have:

Theorem 1.1. *If B is a closed braid with b strands and writhe w , then*

$$(1) \quad \frac{w-b}{2} \leq \hat{g}_{\min}(B) \leq \hat{g}_{\max}(B) \leq \frac{w+b}{2} - 1.$$

If we further assume that the numbers of positive and negative crossings of B are c_+ and c_- , respectively, then

$$(2) \quad (n-1)(w-b) - 2c_- \leq g_{\min}^{(n)}(B) \leq g_{\max}^{(n)}(B) \leq (n-1)(w+b) + 2c_+,$$

Definition 1.2. The braid number $b(L)$ of a link L is the minimal number of strands necessary to represent L as a braid.

Corollary 1.3. *For a link L ,*

$$\begin{aligned} b(L) &\geq \hat{g}_{\max}(L) - \hat{g}_{\min}(L) + 1, \\ b(L) &\geq \frac{1}{2} \limsup_{n \rightarrow \infty} \frac{g_{\max}^{(n)}(L) - g_{\min}^{(n)}(L)}{n-1}. \end{aligned}$$

Proof. $\square$

Theorem 1.4. *If L is a transversal link in the standard contact 3-sphere, then*

$$\begin{aligned} sl(L) &\leq 2\hat{g}_{\min}(L), \\ sl(L) &\leq -\limsup_{n \rightarrow \infty} \frac{g_{\max}^{(n)}(\overline{L})}{n-1}, \end{aligned}$$

where " sl " is the self-linking number, and $\overline{L}$ is the mirror image of L .

Therefore, if K is a Legendrian knot in the standard contact 3-sphere, then

$$\begin{aligned} tb(K) + |r(K)| &\leq 2\hat{g}_{\min}(K), \\ tb(K) + |r(K)| &\leq -\limsup_{n \rightarrow \infty} \frac{g_{\max}^{(n)}(\overline{K})}{n-1}, \end{aligned}$$

where " tb " is the Thurston-Bennequin number, and " r " is the rotation number.

Proof. By [2], any transversal knot is transversally isotopic to a transversal braid, and the self-linking number $sl(B)$ of a transversal braid B with b strands is equal to $w - b$, where w is the writhe of B . So Theorem 1.1 implies the first part of the theorem. The second part follows from the first part by a push-off argument given in [2]. $\square$

In [11], O. Plamenevskaya constructed an invariant ψ of transversal links in standard contact S^3 using the original Khovanov homology $\mathcal{H}$ defined in [6]. We generalize her construction, and, for each $n \geq 2$, define an invariant ψ_n of transversal links in standard contact S^3 using H_n .

Theorem 1.5. *Let L be a transversal link in the standard contact S^3 . For each $n \geq 2$, we can associate to L an element $\psi_n(L)$ of $H_n(\overline{L})$ of cohomological degree 0 and quantum degree $-(n-1)sl(L)$. $\psi_n(L)$ is invariant under transversal isotopy of L up to multiplication by a non-zero scalar.*

When $n = 2$, $\psi_2(L)$ is identified with $\psi(L)$ under the isomorphism

$$H_2(\overline{L}) \cong \mathcal{H}(L) \otimes_{\mathbb{Z}} \mathbb{Q}.$$

Comparing Theorems 1.4 and 1.5, it's easy to see that the sequence $\{\psi_n\}$ sometimes detects the maximal self-linking number. Specially, we can generalize Plamenevskaya's computation about quasi-positive braids, and get a stronger conclusion.

Corollary 1.6. *If L is a transversal link in the standard contact S^3 with $\psi_n(L) \neq 0$ for infinitely many n , then*

$$SL(L) = sl(L) = -\limsup_{n \rightarrow \infty} \frac{g_{max}^{(n)}(\overline{L})}{n-1},$$

where $SL(L)$ is the maximal self-linking number of all transversal links smoothly isotopic to L .

Specially, if a transversal link L is transverse isotopic to a quasi-positive transversal braid, then $\psi_n(L) \neq 0$, $\forall n \geq 2$, and, thus, the above equation is true.

Proof. The first part of the corollary is clear from Theorems 1.4 and 1.5. The second part will be proved in Proposition 4.5. $\square$

A transversal knot invariant is said to be non-classical if it is not determined by the smooth knot type and the self-linking number. There are still no known examples of efficient non-classical transversal knot invariants.

Conjecture 1.7. *The sequence $\{\psi_n\}$ is a non-classical transversal knot invariant.*

The rest of this paper is organized as following: In Section 2, we briefly review the definitions of the Khovanov-Rozansky cohomologies and compare the graded Euler characteristics of these cohomologies with the HOMFLY polynomial, which explains why Theorems 1.1, 1.4 and Corollary 1.3 imply the corresponding results by J. Franks, R. Williams and H. Morton in [4, 9], including the upper bound of the self-linking number from the HOMFLY polynomial. In Section 3, we study the Khovanov-Rozansky cohomologies of closed braids, and prove Theorem 1.1. In Section 4, we construct the invariants ψ_n , and study some of their properties.

2. KHOVANOV-ROZANSKY THEORY

2.1. Matrix factorizations. Let $R = \mathbb{Q}[t_1, \dots, t_m]$ be a polynomial ring. A matrix factorization M over R with potential $w \in R$ is a collection of two R -modules M^0, M^1 and two module homomorphisms $d^0 : M^0 \rightarrow M^1$, $d^1 : M^1 \rightarrow M^0$, called differential maps, s.t.,

$$d^1 d^0 = w \cdot \text{id}_{M^0}, \quad d^0 d^1 = w \cdot \text{id}_{M^1}.$$

We usually write such a matrix factorization M as

$$M^0 \xrightarrow{d_0} M^1 \xrightarrow{d_1} M^0.$$

For $a, b \in R$, denote by $(a, b)_R$ the matrix factorization

$$R \xrightarrow{a} R \xrightarrow{b} R,$$

where a, b act on R by multiplication. This matrix factorization has potential ab . When the polynomial ring R is clear from context, we drop it from the notation, and only write (a, b) .

Denote by

$$\begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \\ \vdots & \vdots \\ a_k & b_k \end{pmatrix}_R$$

the tensor product of $(a_1, b_1)_R, (a_2, b_2)_R, \dots, (a_k, b_k)_R$, where the differential maps are given by the (signed) Leibniz rule. Its potential is

$$w = a_1 b_1 + a_2 b_2 + \dots + a_k b_k.$$

Again, when R is clear from context, we drop it from the notation.

2.2. Grading shifts. Suppose M is a $\mathbb{Z}$ -graded module over a $\mathbb{Z}$ -graded ring, and $m \in \mathbb{Z}$. Define $M\{m\}$ to be M with grading shifted by m , i.e., an element x of $M\{m\}$ is homogeneous of degree $i + m$ if and only if it's a homogeneous element of M of degree i . Similarly, suppose M is a $\mathbb{Z} \oplus \mathbb{Z}$ -graded module over a $\mathbb{Z} \oplus \mathbb{Z}$ -graded ring, and $m, l \in \mathbb{Z}$. Define $M\{m, l\}$ to be M with grading shifted by (m, l) , i.e., an element x of $M\{m, l\}$ is homogeneous of bi-degree $(i + m, j + l)$ if and only if it's a homogeneous element of M of degree (i, j) .

2.3. Planar diagrams. To construct the Khovanov-Rozansky cohomology theories, one considers the planar diagrams Γ with the following properties:

- (1) Γ consists of two types of edges: regular edges and wide edges. These edges intersect only at their endpoints.
- (2) Regular edges are disjoint from each other. Wide edges are disjoint from each other.
- (3) Each regular edge is oriented, and contains at least one marked point. Open endpoints of regular edges are marked.
- (4) Each wide edge has exactly two regular edges entering at one endpoint, and exactly two regular edges exiting from the other endpoint.

See Figures 7, 8, 9, 10 below for examples of such diagrams.

2.4. Matrix factorization associated to a planar diagram. Let Γ be a planar diagram satisfying the conditions in subsection 2.3, and $\{x_1, \dots, x_p\}$ the set of markings on Γ . Let $R' = \mathbb{Q}[x_1, \dots, x_p]$ be the $\mathbb{Z}$ -graded polynomial ring so that $\deg x_i = 2, \forall i$. Let $R = \mathbb{Q}[a, x_1, \dots, x_p]$ be the $\mathbb{Z} \oplus \mathbb{Z}$ -graded polynomial ring so that $\deg a = (2, 0)$ and $\deg x_i = (0, 2), \forall i$.



FIGURE 2. L_j^i

For an oriented (regular) arc L_j^i in Γ from the point marked by x_j to the point marked by x_i that has no marked interior points, let $C_n(L_j^i)$ be the matrix factorization $(\pi_{ij}, x_i - x_j)_{R'}$ given by

$$R' \xrightarrow{\pi_{ij}} R' \{1 - n\} \xrightarrow{x_i - x_j} R',$$

where $\pi_{ij} = x_i^n + x_i^{n-1}x_j + \dots + x_i x_j^{n-1} + x_j^n$. The purpose of the grading shift here is to make $C_n(L_j^i)$ graded in the sense that both π_{ij} and $x_i - x_j$ are homogeneous maps of degree $n + 1$.

Also, let $C(L_j^i)$ be the matrix factorization $(a, x_i - x_j)_R$ given by

$$R \xrightarrow{a} R \{-1, 1\} \xrightarrow{x_i - x_j} R,$$

where the grading shift makes it bi-graded in the sense that a and $x_i - x_j$ are both homogeneous maps of bi-degree $(1, 1)$.




FIGURE 3. Wide edge E

For a wide edge E , let x_i, x_j, x_k, x_l be the closest markings to E as depicted in Figure 3. (It is possible that $x_i = x_k$ or $x_j = x_l$.) Let g be the unique two-variable polynomial such that $g(x + y, xy) = x^{n+1} + y^{n+1}$, and

$$u_1 = u_1(x_i, x_j, x_k, x_l) = \frac{g(x_i + x_j, x_i x_j) - g(x_k + x_l, x_i x_j)}{x_i + x_j - x_k - x_l},$$

$$u_2 = u_2(x_i, x_j, x_k, x_l) = \frac{g(x_k + x_l, x_i x_j) - g(x_k + x_l, x_k x_l)}{x_i x_j - x_k x_l}.$$

Note that u_1 and u_2 are homogeneous polynomials in x_i, x_j, x_k and x_l of degrees $2n$ and $2n - 2$, respectively.

Define $C_n(E)$ to be the matrix factorization

$$\begin{pmatrix} u_1 & x_i + x_j - x_k - x_l \\ u_2 & x_i x_j - x_k x_l \end{pmatrix}_{R'} \{-1\},$$

which is the tensor product of the matrix factorizations

$$R' \{-1\} \xrightarrow{u_1} R' \{-n\} \xrightarrow{x_i + x_j - x_k - x_l} R' \{-1\}$$

and

$$R' \xrightarrow{u_2} R' \{3 - n\} \xrightarrow{x_i x_j - x_k x_l} R'.$$

Again, the grading shifts here make $C_n(E)$ graded in the sense that both differential maps are homogeneous maps of degree $n + 1$.

Also define $C(E)$ to be the matrix factorization

$$\begin{pmatrix} a & x_i + x_j - x_k - x_l \\ 0 & x_i x_j - x_k x_l \end{pmatrix}_R,$$

which is the tensor product of the matrix factorizations

$$R \xrightarrow{a} R \{-1, 1\} \xrightarrow{x_i + x_j - x_k - x_l} R$$

and

$$R \xrightarrow{0} R \{-1, 3\} \xrightarrow{x_i x_j - x_k x_l} R.$$

Again, the grading shifts here make $C(E)$ bi-graded in the sense that both differential maps are homogeneous maps of degree $(1, 1)$.

Finally, we define that

$$C_n(\Gamma) = \left(\bigotimes_{L_j^i} C_n(L_j^i) \right) \bigotimes \left(\bigotimes_E C_n(E) \right),$$

and

$$C(\Gamma) = \left(\bigotimes_{L_j^i} C(L_j^i) \right) \bigotimes \left(\bigotimes_E C(E) \right).$$

where L_j^i runs through all the (regular) oriented arcs starting and ending at marked points with no marked interior points, and E runs through all wide edges.

If Γ is closed, i.e., has no open end points, then $C_n(\Gamma)$ and $C(\Gamma)$ are graded cyclic chain complexes. We denote by $H_n(\Gamma)$ and $H(\Gamma)$ their cohomologies. These cohomologies inherit the gradings of $C_n(\Gamma)$ and $C(\Gamma)$. Note that the $\mathbb{Z}_2$ -grading from the cyclic complex structure is trivial since all the non-vanish cohomology concentrate on one of the two degrees. The cohomologies $H_n(\Gamma)$ and $H(\Gamma)$ do not depend on the choice of marked points.

If Γ has open end points, let $x_{i_1}, \dots, x_{i_q}$ be the markings of these open end points. Then $C_n(\Gamma)$ is a matrix factorization of potential $\sum_{r=1}^q \pm x_{i_r}^{n+1}$, and $C(\Gamma)$ is a factorization of potential $\sum_{r=1}^q \pm a x_{i_r}$, where the sign is "+" if the corresponding open end point is an exit, and is "-" if otherwise. Let I' and I be the ideals generated by $\{x_{i_1}, \dots, x_{i_q}\}$ in R' and R , respectively. Then $C_n(\Gamma)/I'C_n(\Gamma)$ and $C(\Gamma)/IC(\Gamma)$

are graded cyclic chain complex. Again, we denote by $H_n(\Gamma)$ and $H(\Gamma)$ their cohomologies, which have the gradings inherited from $C_n(\Gamma)$ and $C(\Gamma)$. As before, the cohomologies $H_n(\Gamma)$ and $H(\Gamma)$ do not depend on the choice of marked points.




FIGURE 4. Γ_0 and Γ_1

2.5. Khovanov-Rozansky cohomologies of a knot. Let Γ_0 and Γ_1 be the two planar diagrams depicted in Figure 4. Then the matrix factorization $C_n(\Gamma_0)$ is

$$\begin{pmatrix} \pi_{ik} & x_i - x_k \\ \pi_{jl} & x_j - x_l \end{pmatrix}_{R'}.$$

Explicitly, this is

$$\begin{bmatrix} R' \\ R'\{2-2n\} \end{bmatrix} \xrightarrow{P_0^{(n)}} \begin{bmatrix} R'\{1-n\} \\ R'\{1-n\} \end{bmatrix} \xrightarrow{P_1^{(n)}} \begin{bmatrix} R' \\ R'\{2-2n\} \end{bmatrix},$$

where

$$P_0^{(n)} = \begin{pmatrix} \pi_{ik} & x_j - x_l \\ \pi_{jl} & -x_i + x_k \end{pmatrix}, \quad P_1^{(n)} = \begin{pmatrix} x_i - x_k & x_j - x_l \\ \pi_{jl} & -\pi_{ik} \end{pmatrix}.$$

The matrix factorization $C_n(\Gamma_1)$ is

$$\begin{pmatrix} u_1 & x_i + x_j - x_k - x_l \\ u_2 & x_i x_j - x_k x_l \end{pmatrix}_{R'} \{-1\}.$$

Explicitly, this is

$$\begin{bmatrix} R'\{-1\} \\ R'\{3-2n\} \end{bmatrix} \xrightarrow{Q_0^{(n)}} \begin{bmatrix} R'\{-n\} \\ R'\{2-n\} \end{bmatrix} \xrightarrow{Q_1^{(n)}} \begin{bmatrix} R'\{-1\} \\ R'\{3-2n\} \end{bmatrix},$$

where

$$Q_0^{(n)} = \begin{pmatrix} u_1 & x_i x_j - x_k x_l \\ u_2 & -x_i - x_j + x_k + x_l \end{pmatrix}, \quad Q_1^{(n)} = \begin{pmatrix} x_i + x_j - x_k - x_l & x_i x_j - x_k x_l \\ u_2 & -u_1 \end{pmatrix}.$$

Define $\chi_0^{(n)} : C_n(\Gamma_0) \rightarrow C_n(\Gamma_1)$ by the matrices

$$U_0^{(n)} = \begin{pmatrix} x_k - x_j & 0 \\ a_1 & 1 \end{pmatrix}, \quad U_1^{(n)} = \begin{pmatrix} x_k & -x_j \\ -1 & 1 \end{pmatrix},$$

and $\chi_1^{(n)} : C_n(\Gamma_1) \rightarrow C_n(\Gamma_0)$ by the matrices

$$V_0^{(n)} = \begin{pmatrix} 1 & 0 \\ -a_1 & x_k - x_j \end{pmatrix}, \quad V_1^{(n)} = \begin{pmatrix} 1 & x_j \\ 1 & x_k \end{pmatrix},$$

where $a_1 = -u_2 + (u_1 + x_i u_2 - \pi_{jl})/(x_i - x_k)$. These are homomorphisms of matrix factorizations of degree 1, i.e., these commute with the differential maps and raise the

quantum grading by 1. So $\chi_0^{(n)}$ and $\chi_1^{(n)}$ induce homomorphisms of cohomologies. One can check that

$$\chi_1^{(n)} \chi_0^{(n)} = (x_k - x_j) \text{id}_{C_n(\Gamma_0)}, \quad \chi_0^{(n)} \chi_1^{(n)} = (x_k - x_j) \text{id}_{C_n(\Gamma_1)}.$$

There is a similar construction for $C(\Gamma_0)$ and $C(\Gamma_1)$. Explicitly, the matrix factorization $C(\Gamma_0)$ is

$$\begin{bmatrix} R \\ R\{-2, 2\} \end{bmatrix} \xrightarrow{P_0} \begin{bmatrix} R\{-1, 1\} \\ R\{-1, 1\} \end{bmatrix} \xrightarrow{P_1} \begin{bmatrix} R \\ R\{-2, 2\} \end{bmatrix},$$

where

$$P_0 = \begin{pmatrix} a & x_j - x_l \\ a & -x_i + x_k \end{pmatrix}, \quad P_1 = \begin{pmatrix} x_i - x_k & x_j - x_l \\ a & -a \end{pmatrix},$$

and the matrix factorization $C(\Gamma_1)$ is

$$\begin{bmatrix} R \\ R\{-2, 4\} \end{bmatrix} \xrightarrow{Q_0} \begin{bmatrix} R\{-1, 1\} \\ R\{-1, 3\} \end{bmatrix} \xrightarrow{Q_1} \begin{bmatrix} R \\ R\{-2, 4\} \end{bmatrix},$$

where

$$Q_0 = \begin{pmatrix} a & x_i x_j - x_k x_l \\ 0 & -x_i - x_j + x_k + x_l \end{pmatrix}, \quad Q_1 = \begin{pmatrix} x_i + x_j - x_k - x_l & x_i x_j - x_k x_l \\ 0 & -a \end{pmatrix}.$$

Define $\chi_0 : C(\Gamma_0) \rightarrow C(\Gamma_1)$ by the matrices

$$U_0 = \begin{pmatrix} x_k - x_j & 0 \\ 0 & 1 \end{pmatrix}, \quad U_1 = \begin{pmatrix} x_k & -x_j \\ -1 & 1 \end{pmatrix},$$

and $\chi_1 : C(\Gamma_1) \rightarrow C(\Gamma_0)$ by the matrices

$$V_0 = \begin{pmatrix} 1 & 0 \\ 0 & x_k - x_j \end{pmatrix}, \quad V_1^{(n)} = \begin{pmatrix} 1 & x_j \\ 1 & x_k \end{pmatrix}.$$

These are homomorphisms of matrix factorizations, and induce homomorphisms of cohomologies. One can check that χ_0 and χ_1 are of bi-degrees $(0, 2)$ and $(0, 0)$, respectively, and

$$\chi_1 \chi_0 = (x_k - x_j) \text{id}_{C(\Gamma_0)}, \quad \chi_0 \chi_1 = (x_k - x_j) \text{id}_{C(\Gamma_1)}.$$




FIGURE 5. Resolutions of a crossing

Let D be an oriented link diagram. We put markings $\{x_1, \dots, x_m\}$ on D so that none of the crossings is marked, and each arc between two crossings has a marked point. For each crossing in D , there are two ways to resolve it as shown in Figure 5. A resolution of D is a planar diagram obtain from D by resolving all the crossings of D . Together with the markings inherited from D , each resolution of D is a planar diagram satisfying the conditions in Subsection 2.3.

For an arc L_j^i from x_j to x_i that contains no crossings and no other marked points, define $C_n(L_j^i)$ as above, and consider it as the chain complex

$$0 \rightarrow \underbrace{C_n(L_j^i)}_0 \rightarrow 0,$$

where $C_n(L_j^i)$ has cohomological degree 0.

For a positive crossing P_+ in D , let P_0 and P_1 be the two diagrams depicted in Figure 5. Define $C_n(P_+)$ to be the chain complex

$$0 \rightarrow \underbrace{C_n(P_1)\{n\}}_{-1} \xrightarrow{\chi_1^{(n)}} \underbrace{C_n(P_0)\{n-1\}}_0 \rightarrow 0,$$

where $C_n(P_0)\{n-1\}$ has cohomological degree 0, and $C_n(P_1)\{n\}$ has cohomological degree -1 .

For a negative crossing P_- in D , define $C_n(P_-)$ to be the chain complex

$$0 \rightarrow \underbrace{C_n(P_0)\{1-n\}}_0 \xrightarrow{\chi_0^{(n)}} \underbrace{C_n(P_1)\{-n\}}_1 \rightarrow 0,$$

where $C_n(P_0)\{1-n\}$ has cohomological degree 0, and $C_n(P_1)\{-n\}$ has cohomological degree 1.

Now define $C_n(D)$ to be the chain complex

$$C_n(D) = \left(\bigotimes_{L_j^i} C_n(L_j^i) \right) \bigotimes \left(\bigotimes_P C_n(P) \right),$$

where L_j^i runs through all arcs in D starting and ending in marked points that contain no crossings and no other marked points, and P runs through all the crossings of D .

Remark 2.1. (1) Disregarding the chain complex structure on $C_n(D)$, we have

$$C_n(D) = \bigoplus_{\Gamma} C_n(\Gamma)\{p_{\Gamma}\},$$

where Γ runs through all resolutions of D , $p_{\Gamma} \in \mathbb{Z}$. For a given Γ , all elements of $C_n(\Gamma)$ have the same cohomological degree.

- (2) There are two differential maps on $C_n(D)$. One comes from the matrix factorizations $C_n(\Gamma)$, which we denote by d_{mf} . The other comes from the above construction, which we denote by d_{χ} . These two differentials commute.
- (3) $C_n(D)$ is $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}_2$ -graded, where the first $\mathbb{Z}$ -grading is the cohomological grading, the second is inherited from $\mathbb{Z}$ -grading of $C_n(\Gamma)\{p_{\Gamma}\}$, and is called

the quantum grading, and the $\mathbb{Z}_2$ -grading comes from the cyclic chain complex structure of $C_n(\Gamma)$. Note that d_χ preserves the quantum grading.

d_χ induces a differential map on the cohomology $H(C_n(D), d_{mf})$, which we still denote by d_χ . Define $H_n(D)$ to be the cohomology

$$H_n(D) = H(H(C_n(D), d_{mf}), d_\chi).$$

$H_n(D)$ inherits the $\mathbb{Z} \oplus \mathbb{Z}$ -gradings from $C_n(D)$. The induced $\mathbb{Z}_2$ -grading on $H_n(D)$ is trivial since all the non-zero elements concentrate on the degree equal to the $\mathbb{Z}_2$ -number of components of the link represented by D .

M. Khovanov and L. Rozansky proved in [7] that the $\mathbb{Z} \oplus \mathbb{Z}$ -graded isomorphism type of $H_n(D)$ is invariant under Reidemeister moves, and, therefore, is a link invariant. They also showed that, up to multiplication by a non-zero scalar, the homomorphism of H_n induced by a link cobodism is independent of the movie representation. Specially this implies that the H_n theory gives bona fide homology groups, not just isomorphism classes.

The homology $H(D)$ is defined similarly. To L_j^i , P_+ and P_- , we assign the chain complexes

$$\begin{aligned} C(L_j^i) &= [0 \rightarrow \underbrace{C(L_j^i)}_0 \rightarrow 0], \\ C(P_+) &= [0 \rightarrow \underbrace{C(P_1)\{0, -2\}}_0 \xrightarrow{\chi_1} \underbrace{C(P_0)\{0, -2\}}_1 \rightarrow 0], \\ C(P_-) &= [0 \rightarrow \underbrace{C(P_0)\{0, 2\}}_{-1} \xrightarrow{\chi_0} \underbrace{C(P_1)}_0 \rightarrow 0], \end{aligned}$$

where the number below each term indicates its cohomological degree. Then define

$$C(D) = \left(\bigotimes_{L_j^i} C(L_j^i) \right) \otimes \left(\bigotimes_P C(P) \right),$$

where L_j^i runs through all arcs in D starting and ending in marked points that contain no crossings and no other marked points, and P runs through all the crossings of D . Again, we have

$$C(D) = \bigoplus_{\Gamma} C(\Gamma)\{0, q_\Gamma\},$$

where Γ runs through all resolutions of D , $q_\Gamma \in \mathbb{Z}$. Again, there are two differential maps: d_{mf} from the matrix factorization structure, and d_χ from the above chain complex construction. These commute with each other. And d_χ preserves the two quantum gradings. We define

$$H(D) = H(H(C(D), d_{mf}), d_\chi).$$

$H(D)$ inherits the $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ -grading of $C(D)$, i.e. the cohomological grading, the quantum a -grading and the quantum x -grading. M. Khovanov and L. Rozansky proved in [7] that the $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ -graded isomorphism type of $H(D)$ is invariant under Markov moves of braids up to over all shift of grading. Thus, any two braid that represent the

same link have isomorphic H -invariant. Given a braid B with b strands and writhe w , define

$$\hat{H}(B) = H(B) \left\{ \frac{w+b}{2}, \frac{w+b}{2} \right\} \left[-\frac{w+b}{2} \right],$$

where $[-\frac{w+b}{2}]$ means shifting the homological grading by $-\frac{w+b}{2}$. From the construction in [8], it's easy to see that the $\frac{1}{2}\mathbb{Z} \oplus \frac{1}{2}\mathbb{Z} \oplus \frac{1}{2}\mathbb{Z}$ -graded isomorphism type of $\hat{H}(B)$ is honestly invariant under Markov moves of braids.

2.6. Graded Euler characteristics. We normalize the HOMFLY polynomial P by the skein relation

$$\begin{cases} xP(L_-) - x^{-1}P(L_+) = yP(L_0), \\ P(\text{unknot}) = \frac{x-x^{-1}}{y}, \end{cases}$$

where the diagrams of L_+ , L_- and L_0 are identical except at one crossing, where they look like P_+ , P_- and P_0 in Figure 5, respectively.

For a link L , consider the graded Euler characteristic

$$F_n(L) = \sum_{j,k} (-1)^j t^k \dim H_n^{j,k}(L).$$

From [7], we know that F_n is characterized by the following skein relation:

$$\begin{cases} t^n F_n(L_-) - t^{-n} F_n(L_+) = (t - t^{-1}) F_n(L_0), \\ F_n(\text{unknot}) = \frac{t^n - t^{-n}}{t - t^{-1}}. \end{cases}$$

So

$$F_n(L)(t) = P(L)(t^n, t - t^{-1}).$$

Next, consider the graded Euler characteristic

$$\hat{F}(L) = \sum_{j,k,l} (-1)^j t^k q^l \dim \hat{H}^{j,k,l}(L).$$

From [8], we know that $\hat{F}$ is characterized by the following skein relation:

$$\begin{cases} -q^{-1}(-tq)^{\frac{1}{2}} \hat{F}(L_-) + q(-tq)^{-\frac{1}{2}} \hat{F}(L_+) = (q - q^{-1}) \hat{F}(L_0), \\ \hat{F}(\text{unknot}) = -\frac{\sqrt{-1}t^{-\frac{1}{2}}q^{\frac{1}{2}}}{q - q^{-1}}. \end{cases}$$

where L_+ , L_- and L_0 are represented by braid diagrams that identical except at one crossing, where they look like P_+ , P_- and P_0 in Figure 5, respectively.

Change the variables by

$$\begin{cases} x = -q^{-1}(-tq)^{\frac{1}{2}} \\ y = q - q^{-1} \end{cases}$$

Then, in these new variables, we have

$$\begin{cases} x\hat{F}(L_-) - x^{-1}\hat{F}(L_+) = y\hat{F}(L_0), \\ \hat{F}(\text{unknot}) = -x^{-1}y^{-1}. \end{cases}$$

Compare these normalizations with those used in [4, 9]. It's clear that Theorem 1.1 and Corollary 1.3 imply the corresponding results in [4, 9]. Specially, the two upper

bounds of the self-linking number in Theorem 1.4 are sharper than the bound given by the HOMFLY polynomial, which is implicitly proven in [4, 9] (c.f. [3, 5]).

3. RESOLVED BRAIDS

For positive integers b, i with $1 \leq i \leq b-1$, let τ_i be the diagram depicted in Figure 6. That is, from left to right, τ_i consists of $i-1$ upward vertical regular edges, then a vertical wide edge with two regular edges entering through the bottom and two regular edges exiting through the top, and then $b-i-1$ more upward vertical regular edges.




FIGURE 6. τ_i

We use the word $\tau_{i_1} \cdots \tau_{i_m}$ to represent the planar graph formed by stacking the graphs $\tau_{i_1}, \dots, \tau_{i_m}$ together vertically from bottom to top with the top end points of τ_{i_l} identified with the corresponding bottom end points of $\tau_{i_{l+1}}$. The symbol $\tau_{i_1} \cdots \tau_{i_m}$ represents the closed graph obtained from $\tau_{i_1} \cdots \tau_{i_m}$ by attaching a disjoint regular edge from each end point on the top to the corresponding end point at the bottom. We use the convention that the empty word ϕ represents b vertical upward regular edges, and, therefore, ϕ represents b concentric circles. We call $\tau_{i_1} \cdots \tau_{i_m}$ a resolved braid, and $\tau_{i_1} \cdots \tau_{i_m}$ a resolved closed braid. See Figure 7 for examples of such graphs. There are two obvious types of isotopies:

- (I) If $|i-j| > 1$, then $\tau_i \tau_j$ is isotopic to $\tau_j \tau_i$;
- (II) If μ and ν are two words in $\tau_1, \dots, \tau_{b-1}$, then $\underline{\mu\nu}$ is isotopic to $\underline{\nu\mu}$.

We prove inequality (1) in Theorem 1.1 first.

For a word μ in $\tau_1, \dots, \tau_{b-1}$, consider the $\mathbb{Z} \oplus \mathbb{Z}$ -graded cohomology $H(\underline{\mu}) = H(C(\underline{\mu}), d_{mf})$. Denote by $H^{k,l}(\underline{\mu})$ the subspace of $H(\underline{\mu})$ consisting of homogeneous elements of $\mathbb{Z} \oplus \mathbb{Z}$ grading (k, l) , where k is the quantum a -grading, and l is the quantum x -grading. Define


$$g_{\max}(\underline{\mu}) = \max\{k \mid H^{k,l}(\underline{\mu}) \neq 0 \text{ for some } l\},$$

$$g_{\min}(\underline{\mu}) = \min\{k \mid H^{k,l}(\underline{\mu}) \neq 0 \text{ for some } l\}.$$

In order to calculate $H(\underline{\mu})$, we need put marks on regular edges of $\underline{\mu}$. But, as long as there is at least one mark on each regular edge, the result is independent of the choice of the marks.

Lemma 3.1. *For the empty word ϕ (with b strands), we have $g_{\max}(\underline{\phi}) = -1$ and $g_{\min}(\underline{\phi}) = -b$.*

Proof. ϕ represents b concentric circles. We order the circles, and put the marking x_i on the i -th circle. The cyclic Koszul complex $C(\underline{\phi})$ is given over $R = \mathbb{Q}[a, x_1, \dots, x_b]$

FIGURE 7. $\tau_3\tau_2\tau_1$ and $\underline{\tau_3\tau_2\tau_1}$ for $b = 4$.

by

$$\begin{pmatrix} a & 0 \\ \vdots & \vdots \\ a & 0 \end{pmatrix},$$

where there are b rows. After $b - 1$ elementary transformations of rows (c.f. subsection 2.1 of [8]), we see that it is isomorphic to

$$\begin{pmatrix} a & 0 \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix},$$

which is isomorphic to the chain complex

$$C \otimes_{\mathbb{Q}} D^{\otimes(b-1)} \otimes_{\mathbb{Q}} \mathbb{Q}[x_1, \dots, x_b],$$

where C is the chain complex

$$0 \rightarrow \mathbb{Q}[a] \xrightarrow{a} \mathbb{Q}[a]\{-1, 1\} \rightarrow 0,$$

and D is the chain complex

$$0 \rightarrow \mathbb{Q} \xrightarrow{0} \mathbb{Q}\{-1, 1\} \rightarrow 0.$$

Then, by the Künneth Theorem and the Universal Coefficient Theorem, it's easy to check that

$$\begin{aligned} H(\underline{\phi}) &= H(C \otimes_{\mathbb{Q}} D^{\otimes(b-1)} \otimes_{\mathbb{Q}} \mathbb{Q}[x_1, \dots, x_b]) \\ &= H(C) \otimes_{\mathbb{Q}} H(D^{\otimes(b-1)}) \otimes_{\mathbb{Q}} \mathbb{Q}[x_1, \dots, x_b] \\ &= \mathbb{Q}\{-1, 1\} \otimes_{\mathbb{Q}} D^{\otimes(b-1)} \otimes_{\mathbb{Q}} \mathbb{Q}[x_1, \dots, x_b]. \end{aligned}$$

From the definition of D , one can see that the maximal and minimal quantum a -gradings of the right hand side are -1 and $-b$. This proves the lemma. $\square$

Lemma 3.2. *If $\mu = \tau_{i_1} \cdots \tau_{i_m}$ is a word in $\tau_1, \dots, \tau_{b-1}$ with $i_p < i$ for $1 \leq p \leq m$, and $\mu' = \mu\tau_i$, then $g_{\max}(\underline{\mu}) = g_{\max}(\underline{\mu}')$, and $g_{\min}(\underline{\mu}) = g_{\min}(\underline{\mu}')$.*

Proof. The only difference between $\underline{\mu}$ and $\underline{\mu}'$ occurs in the part depicted in Figure 8.



FIGURE 8. $\underline{\mu}$ and $\underline{\mu}'$

The local matrix factorization of the part of $\underline{\mu}$ in Figure 8 is

$$\begin{pmatrix} a & x_1 - x_4 \\ a & 0 \end{pmatrix}.$$

After an elementary transformation, it becomes

$$\begin{pmatrix} a & x_1 - x_4 \\ 0 & 0 \end{pmatrix}.$$

Write $R = \mathbb{Q}[a, x_1, x_2, x_4]$. Explicitly, this matrix factorization is

$$R \oplus R\{-2, 2\} \rightarrow R\{-1, 1\} \oplus R\{-1, 1\} \rightarrow R \oplus R\{-2, 2\},$$

where

$$d^0 = \begin{pmatrix} a & 0 \\ 0 & -x_1 + x_4 \end{pmatrix} \quad \text{and} \quad d^1 = \begin{pmatrix} x_1 - x_4 & 0 \\ 0 & -a \end{pmatrix}.$$

The local matrix factorization of the part of $\underline{\mu}'$ in Figure 8 is

$$\begin{pmatrix} a & x_1 - x_4 \\ 0 & 0 \end{pmatrix}.$$

Explicitly, this matrix factorization is

$$R \oplus R\{-2, 4\} \rightarrow R\{-1, 1\} \oplus R\{-1, 3\} \rightarrow R \oplus R\{-2, 4\},$$

where

$$d^0 = \begin{pmatrix} a & 0 \\ 0 & -x_1 + x_4 \end{pmatrix} \quad \text{and} \quad d^1 = \begin{pmatrix} x_1 - x_4 & 0 \\ 0 & -a \end{pmatrix}.$$

Let f be the map between the two matrix factorizations given by

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} : R \oplus R\{-2, 2\} \rightarrow R \oplus R\{-2, 4\},$$

and

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} : R\{-1, 1\} \oplus R\{-1, 1\} \rightarrow R\{-1, 1\} \oplus R\{-1, 3\}.$$

It's clear that f is an isomorphism of matrix factorizations that preserves the quantum a -grading (but not the x -grading). After tensoring f with the identity maps of other local matrix factorizations of $\underline{\mu}$ (and $\underline{\mu}'$), we get an isomorphism between these matrix factorizations that preserves the a -grading. This proves the lemma. $\square$

Lemma 3.3. *If ν is a word in $\tau_1, \dots, \tau_{b-1}$, $\mu = \nu\tau_i$ and $\mu' = \nu\tau_i\tau_i$, then $g_{\max}(\underline{\mu}) = g_{\max}(\underline{\mu}')$, and $g_{\min}(\underline{\mu}) = g_{\min}(\underline{\mu}')$.*

Proof. (Follow Proposition 30 of [7].) The only difference between $\underline{\mu}$ and $\underline{\mu}'$ occurs in the part depicted in Figure 9.



FIGURE 9. $\underline{\mu}$ and $\underline{\mu}'$

Write $R = \mathbb{Q}[a, x_1, \dots, x_6]$. Let $s_1 = x_5 + x_6$, $s_2 = x_5x_6$, and $\check{R} = \mathbb{Q}[a, x_1, x_2, x_3, x_4, s_1, s_2]$. Then $R = \check{R} \oplus \check{R}\{0, 2\}$.

The local matrix factorization M' of the part of $\underline{\mu}'$ in Figure 8 is

$$\begin{pmatrix} a & x_1 + x_2 - s_1 \\ 0 & x_1x_2 - s_2 \\ a & s_1 - x_3 - x_4 \\ 0 & s_2 - x_3x_4 \end{pmatrix}_R.$$

Let $\check{M}'$ be the matrix factorization over $\check{R}$ given by the same sequence. Then $M' = \check{M}' \oplus \check{M}'\{0, 2\}$. After elementary transformations, $\check{M}'$ is isomorphic to

$$\begin{pmatrix} a & x_1 + x_2 - x_3 - x_4 \\ 0 & x_1x_2 - x_3x_4 \\ 0 & s_1 - x_3 - x_4 \\ 0 & s_2 - x_3x_4 \end{pmatrix}_{\check{R}}.$$

We then exclude the variables s_1, s_2 and remove the last two rows from the above sequence. This does not change chain homotopy type of the matrix factorization

(c.f. Proposition 3 of [8]). The result of this operation is a matrix factorization over $\mathbb{Q}[a, x_1, x_2, x_3, x_4]$ given by the sequence

$$\begin{pmatrix} a & x_1 + x_2 - x_3 - x_4 \\ 0 & x_1 x_2 - x_3 x_4 \end{pmatrix},$$

which is exactly the local matrix factorization of the part of $\underline{\mu}$ depicted in Figure 9. This implies that $H(\underline{\mu}') = H(\underline{\mu}) \oplus H(\underline{\mu})\{0, 2\}$. And the lemma follows. $\square$

Lemma 3.4. *If ν is a word in $\tau_1, \dots, \tau_{b-1}$, $\mu_1 = \nu\tau_{i-1}\tau_i\tau_{i-1}$, $\mu_2 = \nu\tau_i$, and $\mu_3 = \nu\tau_i\tau_{i-1}\tau_i$ then $g_{\max}(\underline{\mu}_3) \leq \max\{g_{\max}(\underline{\mu}_1), g_{\max}(\underline{\mu}_2)\}$, $g_{\min}(\underline{\mu}_3) \geq \min\{g_{\min}(\underline{\mu}_1), g_{\min}(\underline{\mu}_2)\}$.*

Proof. Let $\mu_4 = \nu\tau_{i-1}$. The only difference between $\underline{\mu}_1$, $\underline{\mu}_2$, $\underline{\mu}_3$ and $\underline{\mu}_4$ occurs in the part depicted in Figure 10.




FIGURE 10. $\underline{\mu}_1$, $\underline{\mu}_2$, $\underline{\mu}_3$ and $\underline{\mu}_4$

By Proposition 7 of [8],

$$H(\underline{\mu}_1) \oplus H(\underline{\mu}_2)\{0, 2\} = H(\underline{\mu}_3) \oplus H(\underline{\mu}_4)\{0, 2\}.$$

This implies the lemma. $\square$

Lemma 3.5. *Let $\mu = \tau_{i_1} \dots \tau_{i_m}$ be a word in $\tau_1, \dots, \tau_{b-1}$, such that $m \geq 2$, $i_1 = i_m = i$, and $i_p < i$ for $1 < p < m$. Then, after possibly finitely many steps of isotopies of type (I), there is a sub-word of μ of the form $\tau_j\tau_j$ or $\tau_j\tau_{j-1}\tau_j$ for some $j \leq i$.*

Proof. We induct on the length m of the word μ . When $m = 2$, the lemma is trivially true. Assume that the lemma is true for words with the given properties and length $= 2, \dots, m-1$. Now consider $\mu = \tau_{i_1} \dots \tau_{i_m}$. If there is none or only one τ_{i-1} among $\tau_{i_2}, \dots, \tau_{i_{m-1}}$, then the lemma is true for μ with $j = i$. If there are more than one τ_{i-1} 's among $\tau_{i_2}, \dots, \tau_{i_{m-1}}$, then there is a sub-word $\nu = \tau_{j_1} \dots \tau_{j_l}$ of μ with $2 \leq l < m$, $j_1 = j_l = i-1$, and $j_q < i-1$ for $1 < q < l$. Thus, by induction, the lemma is true for μ . $\square$

Proposition 3.6. *If μ is a word in $\tau_1, \dots, \tau_{b-1}$, then $-b \leq g_{\min}(\underline{\mu}) \leq g_{\max}(\underline{\mu}) \leq -1$.*

Proof. For a word $\mu = \tau_{i_1} \dots \tau_{i_m}$, define the weight $W(\mu)$ of μ to be

$$W(\mu) = \sum_{p=1}^m i_p.$$

We prove the proposition by induction on the weight of a word. If $W(\mu) = 0$, then μ is the empty word ϕ , and Lemma 3.1 implies the proposition. Now assume that the proposition is true for all words with weight $< W$, and $\mu = \tau_{i_1} \cdots \tau_{i_m}$ is a word with weight W . Let $i = \max\{i_1, \dots, i_m\}$. If i appears only once among $i_1, \dots, i_m$, then, after a possible type (II) isotopy, Lemma 3.2 implies that the proposition is true for μ . If i appears at least twice among $i_1, \dots, i_m$, then, after possible type (I) and type (II) isotopies, Lemmas 3.3, 3.4 and 3.5 imply that the proposition is true for μ . $\square$

Proof of inequality (1) in Theorem 1.1. For a braid B with b strands, consider the Khovanov-Rozansky cohomology $H(B)$, and let

$$g_{\max}(B) = \max\{k \mid H^{j,k,l}(L) \neq 0 \text{ for some } j, l\},$$

$$g_{\min}(B) = \min\{k \mid H^{j,k,l}(L) \neq 0 \text{ for some } j, l\}.$$

Note that

$$H(B) = H(H(C(B), d_{mf}), d_\chi),$$

and

$$H(C(B), d_{mf}) = \bigoplus_{\Gamma} H(\Gamma)\{0, q_\Gamma\},$$

where Γ runs through all resolutions of B , and $q_\Gamma \in \mathbb{Z}$. Thus, by Proposition 3.6, we have $-b \leq g_{\min}(B) \leq g_{\max}(B) \leq -1$. But $\hat{H}(B) = H(B)\{\frac{w+b}{2}, \frac{w+b}{2}\}[-\frac{w+b}{2}]$. So

$$\hat{g}_{\max}(B) = g_{\max}(B) + \frac{w+b}{2},$$

$$\hat{g}_{\min}(B) = g_{\min}(B) + \frac{w+b}{2}.$$

And inequality (1) follows. $\square$

The proof of inequality (2) in Theorem 1.1 is similar.

For a word μ in $\tau_1, \dots, \tau_{b-1}$, consider the $\mathbb{Z}$ -graded cohomology $H_n(\underline{\mu}) = H_n(C(\underline{\mu}), d_{mf})$. Denote by $H_n^k(\underline{\mu})$ the subspace of $H_n(\underline{\mu})$ consisting of homogeneous elements of quantum grading k . Define

$$g_{\max}^{(n)}(\underline{\mu}) = \max\{k \mid H_n^k(\underline{\mu}) \neq 0\},$$

$$g_{\min}^{(n)}(\underline{\mu}) = \min\{k \mid H_n^k(\underline{\mu}) \neq 0\}.$$

We have following conclusions similar to Lemmas 3.1, 3.2, 3.3 and 3.4.

Lemma 3.7. (1) *Let ϕ be the empty word (with b strands). Then*

$$g_{\max}^{(n)}(\underline{\phi}) = (n-1)b, \quad \text{and} \quad g_{\min}^{(n)}(\underline{\phi}) = -(n-1)b.$$

(2) *If $\mu = \tau_{i_1} \cdots \tau_{i_m}$ is a word in $\tau_1, \dots, \tau_{b-1}$ with $i_p < i$ for $1 \leq p \leq m$, and $\mu' = \mu\tau_i$, then*

$$g_{\max}^{(n)}(\underline{\mu}') = g_{\max}^{(n)}(\underline{\mu}) - 1, \quad \text{and} \quad g_{\min}^{(n)}(\underline{\mu}') = g_{\min}^{(n)}(\underline{\mu}) + 1.$$

(3) *If ν is a word in $\tau_1, \dots, \tau_{b-1}$, $\mu = \nu\tau_i$ and $\mu' = \nu\tau_i\tau_i$, then*

$$g_{\max}^{(n)}(\underline{\mu}') = g_{\max}^{(n)}(\underline{\mu}) + 1, \quad \text{and} \quad g_{\min}^{(n)}(\underline{\mu}') = g_{\min}^{(n)}(\underline{\mu}) - 1.$$

(4) If ν is a word in $\tau_1, \dots, \tau_{b-1}$, $\mu_1 = \nu\tau_{i-1}\tau_i\tau_{i-1}$, $\mu_2 = \nu\tau_i$, and $\mu_3 = \nu\tau_i\tau_{i-1}\tau_i$ then

$$g_{\max}^{(n)}(\underline{\mu}_3) \leq \max\{g_{\max}^{(n)}(\underline{\mu}_1), g_{\max}^{(n)}(\underline{\mu}_2)\}, \quad \text{and} \quad g_{\min}^{(n)}(\underline{\mu}_3) \geq \min\{g_{\min}^{(n)}(\underline{\mu}_1), g_{\min}^{(n)}(\underline{\mu}_2)\}.$$

Proof. (1) We mark the i -th strand by a single variable x_i of degree 2, for $i = 1, \dots, b$. A straightforward computation shows that

$$H_n(\underline{\phi}) = A_1 \otimes_{\mathbb{Q}} A_2 \otimes_{\mathbb{Q}} \dots \otimes_{\mathbb{Q}} A_b,$$

where $A_i = [\mathbb{Q}[x_i]/(x_i^n)]\{-n+1\}$. The first part of the lemma follows from this.

(2) Note that the $(i+1)$ -th strand in $\underline{\mu}$ is a circle. We mark it by a single variable x of degree 2. Denote by $\hat{\underline{\mu}}$ the closed resolved braid obtained from $\underline{\mu}$ by removing the $(i+1)$ -th strand. Then

$$H_n(\underline{\mu}) = H_n(\hat{\underline{\mu}}) \otimes_{\mathbb{Q}} A = \bigoplus_{i=0}^{n-1} H_n(\hat{\underline{\mu}})\{-n+1+2i\},$$

where $A = [\mathbb{Q}[x]/(x^n)]\{-n+1\}$. But, from Proposition 29 of [7], we know that

$$H_n(\underline{\mu}') = \bigoplus_{i=0}^{n-2} H_n(\hat{\underline{\mu}})\{-n+2+2i\}.$$

Compare these two decompositions. And the second part of the lemma follows.

(3) By Proposition 30 of [7], we have

$$H_n(\underline{\mu}') = H_n(\underline{\mu})\{1\} \oplus H_n(\underline{\mu})\{-1\},$$

which implies the third part of the lemma.

(4) By Proposition 32 of [7], we have

$$H_n(\underline{\mu}_1) \oplus H_n(\underline{\mu}_2) = H_n(\underline{\mu}_3) \oplus H_n(\underline{\mu}_4),$$

where $\mu_4 = \nu\tau_{i-1}$. And the last part of the lemma follows. $\square$

Similar to the proof of Proposition 3.6, we can induct on the weight of a resolved braid using Lemmas 3.5, 3.7, and prove:

Proposition 3.8. *If $\mu = \tau_{i_1} \dots \tau_{i_m}$ is a word in $\tau_1, \dots, \tau_{b-1}$, then*

$$-(n-1)b - m \leq g_{\min}^{(n)}(\underline{\mu}) \leq g_{\max}^{(n)}(\underline{\mu}) \leq (n-1)b + m.$$

Proof. $\square$

Proof of inequality (2) in Theorem 1.1. Let B be a braid with b strands c_+ positive crossings, c_- negative crossings, and writhe $w = c_+ - c_-$. Note that

$$H_n(B) = H(H(C_n(B), d_{mf}), d_\chi),$$

and

$$H(C_n(B), d_{mf}) = \bigoplus_{\Gamma} H_n(\Gamma)\{p_\Gamma\},$$

where Γ runs through all resolutions of B , and $p_\Gamma \in \mathbb{Z}$. Let Γ be any resolution of B . From the construction of $C_n(B)$, it's clear that

$$p_\Gamma = (n-1)w + c_{\Gamma,+} - c_{\Gamma,-},$$

where $c_{\Gamma,+}$ (resp. $c_{\Gamma,-}$) is the number of wide edges in Γ from positive (resp. negative) crossings in B . Let g be the quantum degree of a non-vanishing homogeneous element of $H_n(\Gamma)\{p_\Gamma\}$. Then

$$g_{min}^{(n)}(\Gamma) + p_\Gamma \leq g \leq g_{max}^{(n)}(\Gamma) + p_\Gamma.$$

But, from Proposition 3.8, we have

$$-(n-1)b - c_{\Gamma,+} - c_{\Gamma,-} \leq g_{min}^{(n)}(\Gamma) \leq g_{max}^{(n)}(\Gamma) \leq (n-1)b + c_{\Gamma,+} + c_{\Gamma,-}.$$

So

$$(n-1)(w-b) - 2c_{\Gamma,-} \leq g \leq (n-1)(w+b) + 2c_{\Gamma,+}.$$

It's clear that $c_{\Gamma,+} \leq c_+$, and $c_{\Gamma,-} \leq c_-$. This shows that, if g is the quantum degree of any non-vanishing homogeneous element of $H(C_n(B), d_{mf})$, then

$$(n-1)(w-b) - 2c_- \leq g \leq (n-1)(w+b) + 2c_+.$$

Thus,

$$(n-1)(w-b) - 2c_- \leq g_{min}^{(n)}(B) \leq g_{max}^{(n)}(B) \leq (n-1)(w+b) + 2c_+.$$

□

4. TRANSVERSAL KNOT INVARIANTS $\{\psi_n\}$

In this section, we use the usual notation for braid group. Denote by $\mathcal{B}_b$ the braid group of b strands, and $\sigma_1^{\pm 1}, \dots, \sigma_{b-1}^{\pm 1}$ the standard generators of $\mathcal{B}_b$, i.e., $\sigma_i^{\pm 1}$ permutes the i -th and $(i+1)$ -th strands with a $(\pm)$ -crossing. Also, denote by ϕ the empty word, which is the identity of $\mathcal{B}_b$. Two braids represent the same link if and only if one can be changed into the other by following operations:

- Braid group relations generated by: $\sigma_i \sigma_i^{-1} = \sigma_i^{-1} \sigma_i = \phi$, $\sigma_i \sigma_j = \sigma_j \sigma_i$, when $|i-j| > 1$, and $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$.
- Conjugations: $\mu \leftrightarrow \eta^{-1} \mu \eta$, where $\mu, \eta \in \mathcal{B}_b$.
- Stabilizations and destabilizations:

$$\begin{cases} \text{positive : } \mu (\in \mathcal{B}_b) \leftrightarrow \mu \sigma_b (\in \mathcal{B}_{b+1}) \\ \text{negative : } \mu (\in \mathcal{B}_b) \leftrightarrow \mu \sigma_b^{-1} (\in \mathcal{B}_{b+1}) \end{cases}$$

In [2], Bennequin proved that any transversal link is transversally isotopic to a transversal braid. The following theorem from [10, 12] describes when two transversal braids are transversally isotopic.

Theorem 4.1 ([10, 12]). *If two transversal braids are transversally isotopic, then one can be changed into the other by a finite sequence of braid group relations, conjugations and positive stabilizations and destabilizations.*

Let μ be a closed braid with b strands, and Γ the resolution of μ so that each crossing of μ is 0-resolved as shown in Figure 5. Then Γ consists of b concentric circles, and

$$H_n(\Gamma) = A_1 \otimes_{\mathbb{Q}} A_2 \otimes_{\mathbb{Q}} \cdots \otimes_{\mathbb{Q}} A_b,$$

where $A_i = [\mathbb{Q}[x_i]/(x_i^n)]\{-n+1\}$, and $\deg x_i = 2$ as before. Let w be the writhe of μ . It's clear that the standard isomorphism

$$f : H_n(\Gamma) \xrightarrow{\cong} H_n(\Gamma)\{(n-1)w\} (\subset H(C_n(\mu), d_{mf})),$$

has quantum degree $(n-1)w$, and maps $H_n(\Gamma)$ into cohomological degree 0. Then the element

$$\varphi_n(\mu) = f(x_1^{n-1} \otimes \cdots \otimes x_b^{n-1})$$

has quantum degree $(n-1)(w+b)$ and cohomological degree 0.

Proposition 4.2. $\varphi_n(\mu)$ is a cocycle in the chain complex $(H(C_n(\mu), d_{mf}), d_\chi)$. And, up to multiplication by a non-vanishing scalar, its cohomology class $[\varphi_n(\mu)] \in H_n(\mu) = H(H(C_n(\mu), d_{mf}), d_\chi)$ is invariant under braid group relations, conjugations and negative stabilizations and destabilizations.

Proof. For a negative crossing ς of μ , denote by Γ_ς the resolution of μ obtained by 1-resolving ς and 0-resolving all other crossings. One can easily see that

$$d_\chi(H_n(\Gamma)\{(n-1)w\}) \subset \bigoplus_{\varsigma} H_n(\Gamma_\varsigma)\{(n-1)w-1\} \subset H(C_n(\mu), d_{mf}),$$

where ς runs through all negative crossings of μ . From the second part of Lemma 3.7, we know that the maximal quantum degree of a non-vanishing homogeneous element of $H_n(\Gamma_\varsigma)\{(n-1)w-1\}$ is $(n-1)(w+b)-2$. But d_χ preserves the quantum grading. So $d_\chi(\varphi_n(\mu))$ is a homogeneous element of degree $(n-1)(w+b)$. This implies that $d_\chi(\varphi_n(\mu)) = 0$, and, hence, $\varphi_n(\mu)$ is a cocycle.

Next, we prove the invariance of $[\varphi_n(\mu)]$ under the local moves depicted in Figure 11, which implies the second part of the proposition.




FIGURE 11. Local moves

Remark 4.3. All together, there are 6 braidlike Reidemeister type 3 local moves to deal with. These correspond to the 6 possible ordering of the three branches involved. But each of the other 5 braidlike Reidemeister type 3 local moves is related to move (iii) by braidlike Reidemeister type 2 local moves. So we only need to prove the invariance under move (iii) here.




FIGURE 12. Local diagrams related to move (i)

Invariance under move (i). Consider the planar diagrams in Figure 12.

The local complex associated to μ_0 is

$$0 \rightarrow H_n(\Gamma) \rightarrow 0.$$

And the local complex associated to μ_1 is

$$0 \rightarrow H_n(\Gamma_0)\{1-n\} \rightarrow H_n(\Gamma_1)\{-n\} \rightarrow 0.$$

Note that $H_n(\Gamma_0) = H_n(\Gamma) \otimes A_3$, where $A_3 = [\mathbb{Q}[x_3]/(x_3^n)]\{-n+1\}$, and $\deg x_3 = 2$. There is a homomorphism $A_3 \xrightarrow{\varepsilon} \mathbb{Q}$ of degree $1-n$ given by

$$\varepsilon(x_3^j) = \begin{cases} 1, & \text{if } j = n-1; \\ 0, & \text{otherwise.} \end{cases}$$

This induces a quantum degree preserving homomorphism $\varepsilon : H_n(\Gamma_0)\{1-n\} \rightarrow H_n(\Gamma)$, which, in turn, induces a bi-degree preserving homomorphism of complexes

$$\begin{array}{ccccccc} 0 & \rightarrow & H_n(\Gamma_0)\{1-n\} & \rightarrow & H_n(\Gamma_1)\{-n\} & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & H_n(\Gamma) & \rightarrow & 0 & & \end{array}.$$

It's clear that this chain map maps $\varphi_n(\mu_1)$ to $\varphi_n(\mu_0)$. And, from Section 8 of [7], we know that this chain map induces (a non-zero multiple of) the standard isomorphism from $H_n(\mu_1)$ to $H_n(\mu_0)$. Thus, $[\varphi_n(\mu)]$ is invariant under local move (i) up to multiplication by non-vanishing scalar.

Invariance under moves (ii_a) and (ii_b). The proofs for these two moves are almost identical. So we only prove the invariance under move (ii_a).

Consider the planar diagrams in Figure 13.

The local complex associated to μ_0 is

$$0 \rightarrow H_n(\Gamma_{00}) \rightarrow 0.$$

And the local complex associated to μ_1 is

$$0 \rightarrow H_n(\Gamma_{01})\{1\} \rightarrow H_n(\Gamma_{00}) \oplus H_n(\Gamma_{11}) \rightarrow H_n(\Gamma_{10})\{-1\} \rightarrow 0.$$

Let


$$\pi_{11} : H_n(\Gamma_{00}) \oplus H_n(\Gamma_{11}) \rightarrow H_n(\Gamma_{11})$$

be the natural projection. Consider the quantum degree preserving homomorphisms

$$f_1 = \pi_{11} \circ d_\chi : H_n(\Gamma_{01})\{1\} \rightarrow H_n(\Gamma_{11})$$

and

$$f_2 = d_\chi|_{H_n(\Gamma_{11})} : H_n(\Gamma_{11}) \rightarrow H_n(\Gamma_{10})\{-1\}.$$

FIGURE 13. Local diagrams related to move (ii_a)

Note that

$$H_n(\Gamma_{11}) \cong H_n(\Gamma)\{1\} \oplus H_n(\Gamma)\{-1\}.$$

Let

$$f_{10} : H_n(\Gamma_{01})\{1\} \rightarrow H_n(\Gamma)\{1\} \quad \text{and} \quad f_{21} : H_n(\Gamma)\{-1\} \rightarrow H_n(\Gamma_{10})\{-1\}$$

be the obvious maps induced by f_1 , f_2 and the above isomorphism. Note that Γ_{01} and Γ_{10} are isotopic to Γ . In Section 8 of [7], M. Khovanov and L. Rozansky showed that f_{10} and f_{21} are non-zero multiples of the natural isomorphisms

$$H_n(\Gamma_{01})\{1\} \xrightarrow{\cong} H_n(\Gamma)\{1\} \quad \text{and} \quad H_n(\Gamma)\{-1\} \xrightarrow{\cong} H_n(\Gamma_{10})\{-1\}.$$

Define submodule $A \subset H_n(\Gamma_{00}) \oplus H_n(\Gamma)\{-1\} \subset H_n(\Gamma_{00}) \oplus H_n(\Gamma_{11})$ by

$$A = \{(a, -f_{21}^{-1} \circ d_\chi(a)) \mid a \in H_n(\Gamma_{00})\},$$

and submodule $B \subset H_n(\Gamma_{00}) \oplus H_n(\Gamma_{11})$ by

$$B = d_\chi(H_n(\Gamma_{01})\{1\}).$$

Then

$$H_n(\Gamma_{00}) \oplus H_n(\Gamma_{11}) = A \oplus B \oplus H_n(\Gamma)\{-1\}.$$

It's easy to see that $f_{21} = d_\chi|_{H_n(\Gamma)\{-1\}}$. Thus, the local complex associated to μ_1 splits into the direct sum of the following chain complexes

$$\begin{array}{ccccccc} 0 & & \rightarrow & A & \rightarrow & 0, \\ 0 & \rightarrow & H_n(\Gamma_{01})\{1\} & \xrightarrow{\cong} & B & \rightarrow & 0, \\ & & 0 & \rightarrow & H_n(\Gamma)\{-1\} & \xrightarrow{\cong} & H_n(\Gamma_{10})\{-1\} \rightarrow 0. \end{array}$$

The second and third chain complexes give rise to trivial cohomologies. And the standard isomorphism from $H_n(\mu_0)$ to $H_n(\mu_1)$ is induced by (a non-zero scalar multiple of) the isomorphism of chain complexes

$$F : [0 \rightarrow H_n(\Gamma_{00}) \rightarrow 0] \xrightarrow{\cong} [0 \rightarrow A \rightarrow 0]$$

given by

$$F(a) = (a, -f_{21}^{-1} \circ d_\chi(a)).$$

Since $d_\chi(\varphi_n(\mu_0)) = 0$, it's easy to see that $F(\varphi_n(\mu_0)) = \varphi_n(\mu_1)$. Thus, $[\varphi_n(\mu)]$ is invariant under local move (ii_a) up to multiplication by non-vanishing scalar.

Invariance under moves (iii). The proof in [7] of the invariance of H_n under move (iii) is much more complex than that under moves (i) and (ii). But, if we first use D. Bar-Natan's algebraic trick in [1] to reformulate the description of the isomorphism of cohomologies, then it is quite easy to establish the invariance of $[\varphi_n(\mu)]$ under this move.




FIGURE 14. Local diagrams related to move (iii)

Consider the partially resolved planar diagrams in Figure 14. We extend the definition of the (double) chain complex (C_n, d_{mf}, d_χ) to these diagrams. Then, for $i = 0, 1$, the chain complex $(H(C_n(\mu_i), d_{mf}), d_\chi)$ is the mapping cone of the chain map

$$\chi_{i1} : (H(C_n(\Gamma_{i1}), d_{mf}), d_\chi) \rightarrow (H(C_n(\Gamma_{i0}), d_{mf}), d_\chi),$$

where χ_{i1} is induced by the $\chi_1^{(n)}$ -map associated to the wide edge in Γ_{i1} . Note that Γ_{00} and Γ_{10} are the same diagram, which we denote by Γ . Although formulated differently in [7], M. Khovanov and L. Rozansky actually constructed a chain complex $\mathfrak{C}$ which is a strong deformation retract of both $(H(C_n(\Gamma_{01}), d_{mf}), d_\chi)$ and $(H(C_n(\Gamma_{11}), d_{mf}), d_\chi)$ with corresponding inclusion maps

$$F_i : \mathfrak{C} \rightarrow (H(C_n(\Gamma_{i1}), d_{mf}), d_\chi), \quad i = 0, 1,$$

so that $\chi_{01} \circ F_0 = \chi_{11} \circ F_1$ as homomorphisms from $\mathfrak{C}$ to $(H(C_n(\Gamma), d_{mf}), d_\chi)$. Thus, the following squares commute:

$$\begin{array}{ccccc} (H(C_n(\Gamma_{00}), d_{mf}), d_\chi) & \xrightarrow{\text{id}} & (H(C_n(\Gamma), d_{mf}), d_\chi) & \xleftarrow{\text{id}} & (H(C_n(\Gamma_{10}), d_{mf}), d_\chi) \\ \chi_{01} \uparrow & & \chi \uparrow & & \chi_{11} \uparrow \\ (H(C_n(\Gamma_{01}), d_{mf}), d_\chi) & \xleftarrow{F_0} & \mathfrak{C} & \xrightarrow{F_1} & (H(C_n(\Gamma_{11}), d_{mf}), d_\chi) \end{array},$$

where id is the natural identity homomorphism, and $\chi = \chi_{01} \circ F_0 = \chi_{11} \circ F_1$.

By Lemma 4.5 of [1], for $i = 0, 1$, the maps id and F_i induce an isomorphism $\mathcal{F}_i$ from $H_n(\mu_i)$ to the cohomology of the mapping cones of $\chi : \mathfrak{C} \rightarrow (H(C_n(\Gamma), d_{mf}), d_\chi)$. Then the isomorphism

$$\mathcal{F} = \mathcal{F}_1^{-1} \circ \mathcal{F}_0 : H_n(\mu_0) \rightarrow H_n(\mu_1)$$

is (a multiple of) the standard isomorphism. It's clear that

$$\varphi(\mu_0) \in H(C_n(\Gamma_{00}), d_{mf}) \subset H(C_n(\mu_0), d_{mf}),$$

$$\varphi(\mu_1) \in H(C_n(\Gamma_{10}), d_{mf}) \subset H(C_n(\mu_1), d_{mf}),$$

and

$$\text{id}(\varphi(\mu_0)) = \varphi(\mu_1),$$

where $\text{id} : H(C_n(\Gamma_{00}), d_{mf}) \rightarrow H(C_n(\Gamma_{10}), d_{mf})$ is the natural identity homomorphism. This implies that

$$\mathcal{F}([\varphi(\mu_0)]) = [\varphi(\mu_1)],$$

which shows that $[\varphi_n(\mu)]$ is invariant under local move (iii) up to multiplication by non-vanishing scalar. $\square$

Let L be a transversal link in the standard contact S^3 . We transversally isotope it into a transversal braid.

Definition 4.4. $\psi_n(L) = [\varphi_n(\bar{L})] \in H_n(\bar{L})$.

Proof of Theorem 1.5. From Theorem 4.1 and Proposition 4.2, we know that, up to multiplication by non-zero scalar, $\psi_n(L)$ is independent of the transversal braid representation, and is therefore invariant under transversal isotopy. From the construction of $\psi_n(L) = [\varphi_n(\bar{L})]$, it's clear that it has cohomological degree 0 and quantum degree $(n-1)(-w+b) = -(n-1)sl(L)$, where w is the writhe of L , and b is the number of strands of L . Compare our construction with that by Plamenevskaya in [11]. One easily sees that $\psi_2(L)$ is identified with $\psi(L)$ under the isomorphism $H_2(\bar{L}) \cong \mathcal{H}(L) \otimes_{\mathbb{Z}} \mathbb{Q}$. $\square$

Next we generalize Plamenevskaya's result about quasi-positive transversal braids, which completes the proof of Corollary 1.6.

Proposition 4.5 (Compare Theorem 4 of [11]). *Let $\mu \in \mathcal{B}_b$ be a closed braid, and $\mu' = \mu \sigma_i^{-1}$, where $1 \leq i \leq b-1$. Then there is a homomorphism $f : H_n(\mu') \rightarrow H_n(\mu)$ of quantum degree $n-1$ such that $f([\varphi_n(\mu')]) = [\varphi_n(\mu)]$. Specially, this implies that the ψ_n invariant of a quasi-positive transversal braid is non-vanishing.*

Proof. The local chain complex associated to σ_i^{-1} in $(H_n(\mu', d_{mf}), d_\chi)$ is


$$0 \rightarrow H_n(\Gamma_0)\{1-n\} \rightarrow H_n(\Gamma_1)\{-n\} \rightarrow 0.$$

And the corresponding local complex in $(H_n(\mu, d_{mf}), d_\chi)$ is

$$0 \rightarrow H_n(\Gamma_0) \rightarrow 0.$$

The identity map of $H_n(\Gamma_0)$ induces a local chain map of quantum degree $n-1$:

$$\begin{array}{ccccccc} 0 & \rightarrow & H_n(\Gamma_0)\{1-n\} & \rightarrow & H_n(\Gamma_1)\{-n\} & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & H_n(\Gamma_0) & \rightarrow & 0 & & \end{array}.$$

FIGURE 15. Diagrams Γ_0 and Γ_1

Let $f : H_n(\mu') \rightarrow H_n(\mu)$ be the homomorphism induced by the above local chain map. It's clear that $f([\varphi_n(\mu')]) = [\varphi_n(\mu)]$.

Now let L be a quasi-positive transversal braid of b strands, i.e., L is represented by a braid word of the form

$$\nu = \mu_1 \sigma_{i_1} \mu_1^{-1} \cdots \mu_k \sigma_{i_k} \mu_k^{-1}.$$

Then $\overline{L}$ is represented by the word

$$\overline{\nu} = \overline{\mu}_1 \sigma_{i_1}^{-1} \overline{\mu}_1^{-1} \cdots \overline{\mu}_k \sigma_{i_k}^{-1} \overline{\mu}_k^{-1},$$

where $\overline{\mu}_j$ is the mirror image of μ_j . From the first half of the proposition and the invariance of $[\varphi_n]$, we know that there is a homomorphism $F : H_n(\overline{L}) \rightarrow H_n(\phi)$ such that $F([\varphi_n(\overline{\nu})]) = [\varphi_n(\phi)] \neq 0$, where ϕ is the empty word in b strands. Thus, $\psi_n(L) = [\varphi_n(\overline{\nu})] \neq 0$. $\square$

Some other properties of Plamenevskaya's ψ invariant generalize to ψ_n too. For example:

Proposition 4.6 (Compare Proposition 3 of [11]). *Let $\mu \in \mathcal{B}_b$ be a braid with b strands. If there is an $i \in \{1, \dots, b-1\}$ such that σ_i occurs in the word μ , but σ_i^{-1} does not, then $[\varphi_n(\mu)] = 0$, $\forall n \geq 2$. Specially, if a transversal link L is a transversal stabilization of another transversal link, then $\psi_n(L) = 0$, $\forall n \geq 2$.*

Proof. Fix a crossing σ_i in μ . Let Γ_{-1} be the resolution of μ obtained by 1-resolving σ_i and 0-resolving all other crossings, and Γ_0 the resolution of μ obtained by 0-resolving all the crossings. For a negative crossing ς of μ , let Γ_ς be the resolution of μ obtained by 1-resolving σ_i , ς and 0-resolving all other crossings. Since μ does not contain σ_i^{-1} , Γ_ς is a resolved closed braid of the form $\tau_i \tau_j$, where $j \neq i$. Then, in the chain complex $(H(C_n(\mu), d_{mf}), d_\chi)$, we have

$$d_\chi(H_n(\Gamma_{-1})\{(n-1)w+1\}) \subset H_n(\Gamma_0)\{(n-1)w\} \bigoplus \bigoplus_{\varsigma} H_n(\Gamma_\varsigma)\{(n-1)w\},$$

where w is the writhe of μ , and ς runs through all negative crossings of μ . Let

$$\pi_0 : H_n(\Gamma_0)\{(n-1)w\} \bigoplus \bigoplus_{\varsigma} H_n(\Gamma_\varsigma)\{(n-1)w\} \rightarrow H_n(\Gamma_0)\{(n-1)w\}$$

be the natural projection. Then it's clear that

$$\pi_0 \circ d_\chi = \chi_1 : H_n(\Gamma_{-1})\{(n-1)w+1\} \rightarrow H_n(\Gamma_0)\{(n-1)w\},$$

where χ_1 is induced by the $\chi_1^{(n)}$ -map related to the crossing σ_i . From Proposition 29 of [7], we know that there is a non-zero homogeneous element $\eta \in H_n(\Gamma_{-1})\{(n-1)w+1\}$ of quantum degree $(n-1)(w+b)$ such that $\chi_1(\eta) = \varphi_n(\mu) \in H_n(\Gamma_0)\{(n-1)w\}$. By (1) and (2) of Lemma 3.7, we have that $g_{max}^{(n)}(\Gamma_\varsigma) = (n-1)b-2$ for any negative crossing ς of μ . So the maximal quantum degree of a non-zero homogeneous element of $H_n(\Gamma_\varsigma)\{(n-1)w\}$ is $(n-1)(w+b)-2$. Since d_χ preserves the quantum degree, the projection of $d_\chi(\eta)$ onto $H_n(\Gamma_\varsigma)\{(n-1)w\}$ is zero for any negative crossing ς of μ . This implies that $d_\chi(\eta) = \varphi_n(\mu)$. Thus, $\varphi_n(\mu)$ is a coboundary, and $[\varphi_n(\mu)] = 0$.

If L is a transversal stabilization of another transversal link, then there is a $b \geq 2$, such that L is transversally isotopic to a transversal braid given by a braid word $\mu \in \mathcal{B}_b$, where μ contains one σ_{b-1}^{-1} and no σ_{b-1} . Therefore, $\psi_n(L) = [\varphi_n(\overline{\mu})] = 0$. $\square$

Acknowledgments. The author's interest in finding new bounds for the self-linking number was inspired by Matt Hedden's work on Legendrian knots and the knot Floer homology. The basic idea of Theorem 1.1 was conceived during the Holomorphic Curves Workshop held at IAS in May and June of 2005. The author would like to thank the organizers for their hospitality, and Matt Hedden for interesting conversations. The author would also like to thank the referee for many helpful comments on an earlier draft of this paper.

REFERENCES

- [1] D. Bar-Natan, *Khovanov's homology for tangles and cobordisms*, Algebr. Geom. Topol. **9** (2005), 1443–1499 (electronic).
- [2] D. Bennequin, *Entrelacements et équations de Pfaff*, Astérisque, **107-108** (1983), 87–161.
- [3] E. Ferrand, *On Legendrian knots and polynomial invariants*, Proc. Amer. Math. Soc. **130** (2002), no. 4, 1169–1176 (electronic).
- [4] J. Franks, R. F. Williams, *Braids and the Jones polynomial*, Trans. Amer. Math. Soc. **303** (1987), no. 1, 97–108.
- [5] D. Fuchs, S. Tabachnikov, *Invariants of Legendrian and transverse knots in the standard contact space*, Topology **36** (1997), no. 5, 1025–1053.
- [6] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101** (2000), no. 3, 359–426.
- [7] M. Khovanov, L. Rozansky, *Matrix factorizations and link homology*, arXiv:math.QA/0401268.
- [8] M. Khovanov, L. Rozansky, *Matrix factorizations and link homology II*, arXiv:math.QA/0505056.
- [9] H. Morton, *Seifert circles and knot polynomials*, Math. Proc. Cambridge Philos. Soc. **99** (1986), no. 1, 107–109.
- [10] S. Orevkov, V. Shevchishin, *Markov theorem for transversal links*, J. Knot Theory Ramifications **12** (2003), no. 7, 905–913.
- [11] O. Plamenevskaya, *Transverse knots and Khovanov homology*, arXiv:math.GT/0412184.
- [12] N. Wrinkle, *The Markov Theorem for transverse knots*, arXiv:math.GT/0202055.

DEPARTMENT OF MATHEMATICS AND STATISTICS, LEDERLE GRADUATE RESEARCH TOWER, 710
NORTH PLEASANT STREET, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA 01003-9305, USA
E-mail address: wu@math.umass.edu